maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding an DMB control number.	ion of information. Send comment arters Services, Directorate for Info	s regarding this burden estimate ormation Operations and Reports	or any other aspect of the s, 1215 Jefferson Davis	nis collection of information, Highway, Suite 1204, Arlington
1. REPORT DATE 30 SEP 1997	2 DEDORT TVDE			3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER	
Mapping of Offshore Boundary Layer Structure Using a Scanning Lidar				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory, Remote Sensing Division, Code 7221,4555 Overlook Avenue, Washington, DC, 20375				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAIL Approved for publ	ABILITY STATEMENT ic release; distributi	ion unlimited			
13. SUPPLEMENTARY NO	TES				
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	2	

Report Documentation Page

Form Approved OMB No. 0704-0188

MAPPING OF OFFSHORE BOUNDARY LAYER STRUCTURE USING A SCANNING LIDAR

William Hooper
Remote Sensing Division
Code 7221
Naval Research Laboratory
4555 Overlook Ave
Washington, DC 20375-5351

LONG-TERM GOAL

Develop a rugged, active optical instrument (lidar) that can be used from shore sites, airborne, and ship platforms to characterize the 3-D dynamics of the clear-air boundary layer. Measurements should include not only profiles of temperature, water vapor, the wind vector, and turbulence, but also a 4-D multi-parameter image of the boundary layer.

SCIENTIFIC OBJECTIVES

Measure boundary layer depth, stability of boundary layer, 3-D aerosol structures, and wind field off Wallops Island, Virginia during an ONR-sponsored experiment on shoaling wave clutter propagation scheduled for March 1998. Use the evolution of aerosol structures to statistically measure the wind vector. Also provide 3-D images of coherent boundary layer structures and visualize their 3-D evolution.

From simple single-site profiles, most meteorological models must derive second order flux terms that can be used to derive the change of the temperature, water vapor, and wind fields. The 3-D lidar measurements should provide a direct measurement of key wind variance tensor components that are critical for accurate model development.

APPROACH

During FY-98, the Volume Imaging Lidar (VIL) will be relocated to Wallops Island, Virginia and make measurements as part of the planned field experiment. As part of this experiment, RHI, PPI and specialized scans will be made. Boundary layer depth, stability, and wind fields will be derived from these data.

WORK COMPLETED

During FY-97, lidar was upgraded, tested, and calibrated in preparation for field experiment at Wallops Island, Virginia.

RESULTS

FY-97 is the first year of funding. Results will be reported after the Wallops Island experiment.

IMPACT/APPLICATION

FY-97 is the first year of funding. Expect results from Wallops Island experiment to improve understanding of Coastal Boundary Layer and air-sea interaction.

TRANSITIONS

None, FY-97 is the first year of funding.

RELATED PROJECTS

As part of ONR funded research at NRL, the lidar is also being used to (1) characterize the generation and processing of aerosol in coastal regions and (2) develop techniques for measuring optical refraction.

REFERENCES

- W. P. Hooper and E. W. Eloranta, 1986. "Lidar Measurements of Wind in the Planetary Boundary Layer: The Method, Accuracy, and Results from Joint Measurements with Radiosonde and Kytoon," *J. Cli. and Appl. Meteor.* 25, 990-1001.
- J. E. James and W. P. Hooper, 1995. "NRL Volume Imaging Lidar and Its Capabilities," NRL/MR/7221-95-7678.
- W. P. Hooper, J.E. James, and R. J. Lind, 1996. "Lidar Observations of Turbulent Vortex Shedding by an Isolated Topographic Feature," *Bound. Layer Meteor.* 80, 95-108.